

Safina Ali and Daniel J. Drucker

Am J Physiol Endocrinol Metab 296:415-421, 2009. First published Dec 30, 2008;
doi:10.1152/ajpendo.90887.2008

You might find this additional information useful...

This article cites 76 articles, 26 of which you can access free at:

<http://ajpendo.physiology.org/cgi/content/full/296/3/E415#BIBL>

Updated information and services including high-resolution figures, can be found at:

<http://ajpendo.physiology.org/cgi/content/full/296/3/E415>

Additional material and information about *AJP - Endocrinology and Metabolism* can be found at:

<http://www.the-aps.org/publications/ajpendo>

This information is current as of April 6, 2009 .

Benefits and limitations of reducing glucagon action for the treatment of type 2 diabetes

Safina Ali and Daniel J. Drucker

Department of Medicine, Samuel Lunenfeld Research Institute, Mt. Sinai Hospital, Toronto, Ontario, Canada

Submitted 22 February 2008; accepted in final form 24 December 2008

Ali S, Drucker DJ. Benefits and limitations of reducing glucagon action for the treatment of type 2 diabetes. *Am J Physiol Endocrinol Metab* 296: E415–E421, 2009. First published December 30, 2008; doi:10.1152/ajpendo.90887.2008.— Glucagon is secreted from the α -cells of the pancreatic islets and regulates glucose homeostasis through modulation of hepatic glucose production. As elevated glucagon levels contribute to the pathophysiology of hyperglycemia in subjects with type 2 diabetes, reduction of glucagon receptor gene (*Gcgr*) activity represents a potential target for the treatment of T2DM. Herein, we review current concepts of glucagon action in hepatic and extrahepatic tissues and evaluate the therapeutic potential, mechanisms of action, and safety of reducing *Gcgr* signaling for the treatment of T2DM.

INSULIN AND GLUCAGON ARE PANCREATIC HORMONES that play pivotal roles in regulating glucose homeostasis and metabolism. Subjects with type 2 diabetes mellitus (T2DM) have fasting hyperglucagonemia with impaired postprandial glucagon suppression in the presence of defective insulin secretion and/or action. Inhibition of postprandial glucagon secretion in diabetic subjects has been shown to substantially reduce blood glucose, suggesting that glucagon contributes significantly to the hyperglycemia seen in subjects with T2DM (16, 62, 63). These observations have prompted interest in reduction of glucagon secretion or blockade of glucagon receptor gene (*Gcgr*) activity for the treatment of T2DM. In this review, we summarize recent advances in understanding the actions of glucagon in hepatic and extrahepatic tissues and provide an overview of the therapeutic potential of *Gcgr* inhibition for the treatment of T2DM.

Glucagon Synthesis and Secretion

Glucagon is a 29-amino acid peptide hormone encoded within a proglucagon precursor that also contains amino acid sequences for glucagon-like peptide (GLP)-1, GLP-2, oxyntomodulin, and glicentin (Fig. 1). Tissue-specific posttranslational processing of proglucagon in α -cells is mediated by prohormone convertase (PC)2, which cleaves proglucagon to liberate glucagon and leaves unprocessed the carboxy-terminal major proglucagon fragment. In contrast, PC1/3 cleaves proglucagon in intestinal L cells and brain to liberate GLP-1, GLP-2, oxyntomodulin, and glicentin (3). Glucagon secretion by α -cells is highly regulated by multiple factors, the most important of which are glucose and insulin (18) and are reviewed in Ref. 61. Low glucose levels activate specific channels in the brain, specifically the ATP-sensitive K^+ (KATP) channel (51), and on pancreatic α -cells to generate action potentials of sodium and calcium currents, leading to

glucagon secretion. However, whether the modulating effect of glucose on glucagon secretion is predominantly direct or indirect remains uncertain. Studies conducted with mouse and human α -cells show that glucose can directly inhibit glucagon secretion. In contrast, studies with rat α -cells show that glucose regulates glucagon secretion in a paracrine manner (61). β -Cell-derived products such as insulin, GABA, and zinc also inhibit glucagon secretion (5). The mechanisms responsible for insulin-mediated inhibition of α -cell glucagon secretion may involve insulin-mediated activation of GABA receptor translocation to the cell surface in an Akt-dependent manner (74). Similarly, secretion of zinc from β -cells appears to be important for suppression of glucagon secretion, and reduced zinc secretion promotes enhanced glucagon secretion in response to hypoglycemia (76). Nevertheless, experiments using rodent and human islets demonstrated that glucose-mediated suppression of glucagon secretion may occur independently of GABA or zinc and requires functional KATP channels (48). Somatostatin inhibits glucagon secretion by inhibition of adenylate cyclase and cAMP production, and genetic deletion of the somatostatin receptor subtype 2 is associated with mild hyperglucagonemia and defective glucose- and somatostatin-mediated suppression of glucagon secretion in isolated islets *in vitro* (65). Similarly, the incretin hormone GLP-1 inhibits glucagon secretion in a glucose-dependent manner through mechanisms requiring the somatostatin receptor subtype 2 (15).

Glucagon Action and the *Gcgr*

The major biological action of glucagon is to counteract the actions of insulin and maintain normoglycemia during the fasting state by inducing hepatic glucose production. Glucagon exerts its action on target tissues through activation of the *Gcgr*, a G protein-coupled receptor member of the class II G protein-coupled receptor superfamily (33). *Gcgr* activation leads to signal transduction by G proteins ($G_{s\alpha}$ and G_q), whereby $G_{s\alpha}$ activates adenylate cyclase, which causes cAMP production, resulting in an increase in levels of protein kinase A. G_q activation leads to phospholipase C-mediated increases

Address for reprint requests and other correspondence: D. J. Drucker, Mt. Sinai Hospital, 60 Murray St., Mailbox 39, Toronto, ON, Canada M5T 3L9 (d.drucker@utoronto.ca).




Fig. 1. A: structures of proglucagon B: tissue-specific posttranslational processing of proglucagon in the pancreas gives glicentin-related polypeptide (GRPP), glucagon, intervening peptide-1 (IP-1), and major proglucagon fragment (MPGF), whereas glicentin, oxyntomodulin (OXM), intervening peptide (IP-2), glucagon-like peptide (GLP)-1, and GLP-2 are liberated from the brain and the intestine. S, signal peptide.

in intracellular calcium levels. Gcgr signaling in the liver results in increased hepatic glucose production by induction of glycogenolysis and gluconeogenesis along with inhibition of glycogenesis (34). The actions of glucagon to promote increased hepatic glucose production are extremely rapid and reflect changes in the activity of enzymes regulating gluconeogenesis and glycogenolysis. Glucagon-stimulated increases in cAMP lead to activation of glycogen phosphorylase and inhibition of glycogen synthase. The actions of glucagon to control gluconeogenesis are mediated through coordinate regulation of the cAMP-regulated binding protein, regulated transcription coactivator 2, histone acetyltransferase p300, and the nutrient-sensing deacetylase sirtuin 1, resulting in increased expression of genes regulating gluconeogenesis (45). The Gcgr is also expressed in extrahepatic tissues, which includes heart, intestinal smooth muscle, kidney, brain, and adipose tissue (27), and much less is known about the action of glucagon in these tissues (Fig. 2).

Glucagon and the Cardiovascular System

Pharmacological doses of glucagon activate adenylate cyclase in a β -adrenoreceptor-independent manner, leading to cAMP production in the myocardium and a positive inotropic and chronotropic effect. Accordingly, glucagon is occasionally used for the treatment of poisoning caused by cardiodepressant drugs such as β -blockers or calcium channel blockers (73). The inotropic effects mediated by glucagon in the cardiovascular system may be preferentially localized to the ventricular myocardium due in part to differential Gcgr expression in the ventricle compared with the atrium (23). Infusion of glucagon in perfused rat hearts at levels designed to achieve physiolog-

ical concentrations of glucagon leads to induction of glycolysis and glucose oxidation, similar to insulin actions in the heart that are mediated via phosphatidylinositol 3-kinase-dependent and adenylate cyclase- and cAMP-independent pathways (28). Hence, unlike the effects of glucagon in the liver that generally oppose insulin action, glucagon and insulin action in the heart may overlap in regard to stimulation of fuel metabolism.

Glucagon, the Kidney, and the Gastrointestinal Tract

Glucagon stimulates adenylate cyclase and cAMP production in nephrons and in cell-free preparations of human renal medullas (54, 75). Although the role of glucagon in the control of renal glucose output remains uncertain, glucagon regulates the rate of kidney filtration, urea excretion, and water reabsorption by the kidney (26) via direct and indirect mechanisms (4). Paradoxically, long-term infusion of glucagon in mice leads to kidney injury through the development of hypertension, hypertrophy, and increased proliferation of mesangial cells (40). Although the Gcgr is also expressed in the gut, where it regulates motility (58, 69), very little is known about the physiological role of glucagon in the gut.

Glucagon and the Endocrine Pancreas

Gcgr immunoreactivity and mRNA expression have been detected predominantly in β -cells from rodent pancreas; however, subsets of α - and δ -cells also express the Gcgr (37). Additionally, glucagon has been shown to regulate cAMP production in β -cells. However, glucagon-mediated cAMP production in β -cells is less potent than that induced by the incretin hormones GLP-1 and gastric inhibitory polypeptide (GIP) (53). Nevertheless, glucagon induces insulin secretion in human subjects. Moreover, insulin secretion is increased from perfused pancreas and isolated β -cells in the presence of glucagon (37, 53). The stimulatory actions of glucagon on the islet β -cell may be mediated through dual activation of both the Gcgr and the GLP-1 receptor (GLP-1R) (52). However, the molecular mechanism(s) and physiological importance of glucagon-stimulated insulin secretion require further elucidation. Even less is known about the role of the Gcgr in α -cells; however, several studies have demonstrated Gcgr expression in at least a subset of rodent α -cells (37, 47). Glucagon stimulates cAMP production in a dose-dependent manner from rat and mouse α -cells (37, 47) and increases α -cell exocytosis in a PKA-dependent manner, suggesting that it may regulate its own secretion (47). However, the importance of glucagon action on α -cells is uncertain.

Glucagon Action in the Brain and Adipose Tissue

The proglucagon gene is expressed in the brainstem and, to a lesser extent, in the hypothalamus (17), and afferent projections distribute proglucagon-derived peptides to diverse brain regions (35). Glucagon binds to brain membranes and mouse astrocytes and stimulates adenylate cyclase and cAMP production, respectively (12, 31). Intracerebral administration of glucagon at doses likely to produce pharmacological levels of glucagon in the brain produces dose-dependent hyperglycemia in rodents through mechanisms requiring cholinergic and α -adrenergic neural pathways (1, 50). Glucagon infusion in the central nervous system also inhibits food intake (32), and the anorectic actions of glucagon require functional vagal afferents (72). Moreover, neutralization




Fig. 2. Glucagon action in hepatic and extrahepatic tissues. The actions of glucagon are well studied in the liver (solid gray arrow). The actions of glucagon in extrahepatic tissues, including kidney, gastrointestinal tract, pancreas, adipose tissue, cardiovascular system, and the central nervous system, are less extensively defined (dashed gray arrow).

of endogenous glucagon via intraportal infusion of glucagon antibodies increased meal size in normal rats, effects that were abolished in rats with selective hepatic vagotomy (20). The satiety-promoting effects of glucagon may also involve suppression of ghrelin secretion, actions that require an intact hypothalamic-pituitary axis, and ghrelin has been shown to regulate feeding behavior, suggesting that the satiety effect of glucagon could be mediated through ghrelin (2).

Although Gcgr expression is detected in adipose tissue, the role of the Gcgr in the induction of lipolysis in adipose tissue is contradictory. Although glucagon increases lipolysis in rat and human adipocytes (29, 59), subcutaneous infusion of glucagon in abdominal adipose tissue of healthy male subjects had no effect on lipolysis (24). Hence, the precise role of glucagon in the control of lipolysis remains uncertain.

Glucagon and the Pathophysiology of Type 2 Diabetes

Type 2 diabetes is characterized by impaired insulin secretion and/or action, and many subjects also exhibit inappropriate levels of

circulating glucagon in the fasting and postprandial state. An increase in the glucagon/insulin ratio is likely an important determinant of the hyperglycemia seen in type 2 diabetes patients (6, 7, 19). Consistent with the importance of glucagon for fasting hyperglycemia, infusion of low doses of glucagon leads to the development of hyperglycemia (44), whereas suppression of glucagon secretion in the fasting state by somatostatin infusion significantly reduces hepatic glucose production (6). Lack of suppression of postprandial glucagon secretion in subjects with T2DM also plays an important role in the pathogenesis of postprandial hyperglycemia (22, 30, 63). The molecular mechanisms responsible for dysregulation of α-cell glucagon secretion in diabetic subjects remain unclear but may include impaired glucose sensing by α-cells and/or resistance of α-cells to the inhibitory actions of insulin or other β-cell secretory products such as zinc or GABA.

Reduction of Gcgr Signaling for the Treatment of Diabetes

Considerable preclinical evidence supports the targeting of glucagon action as an effective approach to reduction of hyperglycemia. Immunoneutralization of glucagon with a

monoclonal antibody produced significant improvements in plasma glucose in rats with streptozotocin-induced diabetes (10). Similarly, glucagon antibodies markedly reduced hepatic glucose production and reduced the extent of hyperglycemia in normal and diabetic rabbits (9). Additionally, immunoneutralization of plasma glucagon decreased hepatic glucose output and reduced glucose and Hb A_{1c} in *ob/ob* mice, providing further evidence for the central role of glucagon in the pathogenesis of diabetic hyperglycemia (67).

Both peptide and nonpeptide glucagon receptor antagonists have been generated for use as experimental tools to block glucagon action (34). Consistent with data from glucagon immunoneutralization studies, *Gcgr* antagonists lower blood glucose in response to exogenous glucagon administration in nondiabetic rodents and block the actions of endogenously elevated levels of glucagon, leading to reduction of hyperglycemia in diabetic rodents (14, 36, 70). Several different classes of small molecule-based orally available *Gcgr* antagonists have been identified, including trisubstituted ureas, benzimidazole, alkylidene hydrazides, and β -alanine derivatives. These molecules were active following oral administration in dogs, rhesus monkeys, and nondiabetic and diabetic rodents (38, 39, 42, 49). Furthermore, BAY27-995, a small-molecule *Gcgr* antagonist, successfully blocked exogenous glucagon-stimulated glucose production in human subjects (60).

Complementary strategies for reduction of hepatic *Gcgr* signaling have utilized antisense oligonucleotide (ASO) to target hepatic *Gcgr* expression. Twice weekly intraperitoneal administration of *Gcgr* ASOs to *db/db* mice significantly reduced plasma levels of glucose, triglycerides, and free fatty acids without associated hypoglycemia (43). Similarly, *Gcgr* ASOs reduced hyperglycemia in *ob/ob* and *db/db* mice and Zucker diabetic fatty rats together with a reduction in plasma and hepatic triglyceride content. Intriguingly, plasma levels of glucagon and GLP-1 were markedly elevated in rodents treated with *Gcgr* ASOs, in association with the development of α -cell hyperplasia and hypertrophy, findings that were reversible following discontinuation of ASO therapy (66). Taken together, these studies demonstrate that transient inhibition of *Gcgr* expression and/or glucagon action can inhibit hepatic glucose production, leading to improved glucose homeostasis in rodents.

Elimination of *Gcgr* Signaling: Insights From *Gcgr*^{-/-} Mice

Studies of mice with targeted disruption of the *Gcgr* gene (*Gcgr*^{-/-}) have demonstrated that *Gcgr*^{-/-} mice are viable and exhibit mild fasting hypoglycemia (21, 56). Unexpectedly, *Gcgr*^{-/-} mice exhibit markedly increased circulating levels of GLP-1 and multiple phenotypes consistent with enhanced GLP-1 action, including improved glucose homeostasis, reduced gastric emptying, decreased adiposity, increased lean body mass, and resistance to streptozotocin-induced diabetes and diet-induced obesity (Table 1) (13, 21, 56, 68, 71). The improvement in glucose homeostasis likely reflects a reduction in fasting glycemia due to reduced hepatic glucose production and improved β -cell function as a result of increased circulating levels of GLP-1 in *Gcgr*^{-/-} mice (21). Hence, the extent to which the improved metabolic phenotype of *Gcgr*^{-/-} mice reflects the direct loss of *Gcgr* signaling in liver vs. the contribution of enhanced GLP-1 action remains to be deter-

Table 1. Comparison of phenotypes in *Gcgr*^{-/-} (increased GLP-1 action) vs. *Glp1r*^{-/-} mice (lack of GLP-1 action)

| | <i>Gcgr</i> ^{-/-} (\uparrow GLP-1 action) | <i>Glp1r</i> ^{-/-} (\downarrow GLP-1 action) |
|--|--|---|
| Fasting glycemia | Hypoglycemia | Hyperglycemia |
| Oral and intraperitoneal glucose tolerance | Improved | Impaired |
| Susceptibility to STZ-induced diabetes | Decreased | Increased |
| Gastric emptying | Reduced | Normal |

Gcgr, glucagon receptor gene; GLP-1, glucagon-like peptide-1; *Glp1r*, GLP-1 receptor; STZ, streptozotocin.

mined through additional studies employing GLP-1 receptor antagonists and/or genetic loss of GLP-1 action in the context of reduced *Gcgr* signaling.

Inhibition of *Gcgr* Signaling: Potential Limitations

Although inhibition of *Gcgr* signaling for the treatment of T2DM shows promising results in rodent models with diabetes, partial or complete ablation of the *Gcgr* is associated with several unexpected phenotypes that merit careful consideration in light of therapeutic attempts to attenuate *Gcgr* action for the treatment of type 2 diabetes. Notably, loss of *Gcgr* signaling is associated with the development of islet hyperplasia and increased endocrine cell proliferation, detectable in rodents following partial transient reduction or genetic extinction of hepatic *Gcgr* signaling (21, 66). The long-term safety of activating signaling pathways that promote increased islet cell proliferation has not been defined; however, it is worth noting that rodent β -cells tend to exhibit a greater capacity for replication relative to human β -cells (57). Moreover, *Gcgr*^{-/-} mice exhibit significant increases in pancreatic weight (21), likely reflecting changes in cell number within the exocrine pancreas, although a precise analysis of the cellular composition of the *Gcgr*^{-/-} pancreas has not been forthcoming.

More recent studies have focused on the function and viability of *Gcgr*^{-/-} hepatocytes. Exogenous administration of glucagon induces a hypolipidemic effect in multiple species (8, 25), and glucagon administration promotes mobilization of hepatic fat in lactating dairy cows (55). Consistent with these findings, fasted *Gcgr*^{-/-} mice exhibit significant defects in lipid synthesis, secretion, and oxidation (46) and fail to up-regulate a gene expression program promoting lipid oxidation. Moreover, glucagon exerts multiple hypolipidemic actions directly on hepatocytes, in part through a PPAR α -dependent pathway. Furthermore, high-fat feeding of *Gcgr*^{-/-} mice was associated with accelerated development of steatosis in some (46) but not all studies (13). These findings imply that a threshold level of *Gcgr* signaling is required for hepatocytes to regulate synthesis, secretion, and oxidation of lipids, and marked attenuation of *Gcgr* signaling would be predicted to be associated with an increased risk of dyslipidemia and fatty liver.

Complementary studies have examined the role of the *Gcgr* as a determinant of hepatocyte viability. The class 2 family of G protein-coupled receptors contains several members, notably the GIP receptor, GLP-1R, and GLP-2R, that have been linked to control of cell survival. For example, increased GLP-1R signaling reduces β -cell death both in cell culture studies in

vitro and in multiple preclinical studies in vivo (11, 41), whereas genetic disruption of the *Glp1r* is associated with enhanced susceptibility to apoptotic injury (41). Sinclair et al. (64) demonstrated that glucagon administration exerted cytoprotective actions for hepatocytes cultured in vitro or following glucagon administration to mice in vivo. Conversely, *Gcgr*^{-/-} mice exhibit significantly enhanced susceptibility to experimental liver injury either following exogenous administration of the proapoptotic Fas ligand or after high-fat feeding (64). Moreover, reintroduction of the *Gcgr* in *Gcgr*^{-/-} mice by adenoviral gene transfer significantly attenuated the development of liver injury in vivo. These findings demonstrate that the *Gcgr* is an important regulator of hepatocellular survival; however, the minimum level of *Gcgr* expression required for optimization of hepatocyte survival has not been determined.

Another aspect of *Gcgr* biology that requires additional attention is the ability of hepatocytes with reduced *Gcgr* signaling to mount an appropriate counterregulatory response to hypoglycemia. Unexpectedly, genetic elimination of the *Gcgr* augments the counterregulatory response to hypoglycemia in *Gcgr*^{-/-} mice (21). The finding that *Gcgr*^{-/-} mice exhibit increased epinephrine-stimulated cAMP production in liver membranes may contribute to the maintenance of an appropriate counterregulatory response in the absence of glucagon action. Nevertheless, more detailed studies are required that examine the relationship between reduced *Gcgr* signaling and the counterregulatory response to hypoglycemia in human subjects.

Summary

The central importance of glucagon action for regulation of hepatic glucose production, taken together with considerable preclinical and clinical evidence documenting the contribution of dysregulated glucagon secretion to the pathophysiology of diabetic hyperglycemia, makes the *Gcgr* a logical target for the treatment of T2DM. Moreover, compelling data from preclinical studies illustrates the therapeutic potential of *Gcgr* antagonists or molecules targeting the expression of the *Gcgr*. Nevertheless, the *Gcgr*^{-/-} mouse exhibits several unexpected phenotypes, namely α -cell hyperplasia and increased mass of the pancreas, that complicate ascertainment of the risk/benefit ratio for marked inhibition of *Gcgr* signaling. Moreover, the demonstration that complete reduction of *Gcgr* signaling increases the susceptibility to hepatosteatosis and hepatocellular injury raises further questions about the margin for safety in reduction of glucagon action. Furthermore, the central role of glucagon as the primary hormone responsible for the counterregulatory response to hypoglycemia requires a critical evaluation of the extent to which reduction of glucagon action can be safely achieved in diabetic subjects treated with agents that may inadvertently lead to the development of hypoglycemia. Future studies in normal and diabetic human subjects should identify the extent to which reduction of *Gcgr* signaling produces a compelling therapeutic benefit without incurring a risk of adverse events.

GRANTS

The Canadian Institute of Health Research provided an operating grant; D. Drucker is also supported by a Canada Research Chair in Regulatory Peptides. S. Ali was supported by a Meredith and Malcolm Silver Cardiovascular Award (graduate studentship award).

REFERENCES

1. Amir S. Central glucagon-induced hyperglycemia is mediated by combined activation of the adrenal medulla and sympathetic nerve endings. *Physiol Behav* 37: 563–566, 1986.
2. Arafat AM, Perschel FH, Otto B, Weickert MO, Rochlitz H, Schoff C, Spranger J, Mohlig M, Pfeiffer AF. Glucagon suppression of ghrelin secretion is exerted at hypothalamus-pituitary level. *J Clin Endocrinol Metab* 91: 3528–3533, 2006.
3. Baggio LL, Drucker DJ. Biology of incretins: GLP-1 and GIP. *Gastroenterology* 132: 2131–2157, 2007.
4. Bankir L, Ahloulay M, Devreotes PN, Parent CA. Extracellular cAMP inhibits proximal reabsorption: are plasma membrane cAMP receptors involved? *Am J Physiol Renal Physiol* 282: F376–F392, 2002.
5. Bansal P, Wang Q. Insulin as a physiological modulator of glucagon secretion. *Am J Physiol Endocrinol Metab* 295: E751–E761, 2008.
6. Baron AD, Schaeffer L, Shragg P, Kolterman OG. Role of hyperglucagonemia in maintenance of increased rates of hepatic glucose output in type II diabetics. *Diabetes* 36: 274–283, 1987.
7. Basu A, Shah P, Nielsen M, Basu R, Rizza RA. Effects of type 2 diabetes on the regulation of hepatic glucose metabolism. *J Investig Med* 52: 366–374, 2004.
8. Bobe G, Ametaj BN, Young JW, Beitz DC. Effects of exogenous glucagon on lipids in lipoproteins and liver of lactating dairy cows. *J Dairy Sci* 86: 2895–2903, 2003.
9. Brand CL, Jorgensen PN, Svendsen I, Holst JJ. Evidence for a major role for glucagon in regulation of plasma glucose in conscious, nondiabetic, and alloxan-induced diabetic rabbits. *Diabetes* 45: 1076–1083, 1996.
10. Brand CL, Rolin B, Jorgensen PN, Svendsen I, Kristensen JS, Holst JJ. Immunoneutralization of endogenous glucagon with monoclonal glucagon antibody normalizes hyperglycaemia in moderately streptozotocin-diabetic rats. *Diabetologia* 37: 985–993, 1994.
11. Brubaker PL, Drucker DJ. Glucagon-like peptides regulate cell proliferation and apoptosis in the pancreas, gut and central nervous system. *Endocrinology* 145: 2653–2659, 2004.
12. Cockram CS, Kum W, Ho SK, Zhu SO, Young JD. Binding and action of glucagon in cultured mouse astrocytes. *Glia* 13: 141–146, 1995.
13. Conarello SL, Jiang G, Mu J, Li Z, Woods J, Zycband E, Ronan J, Liu F, Roy RS, Zhu L, Charron MJ, Zhang BB. Glucagon receptor knockout mice are resistant to diet-induced obesity and streptozotocin-mediated beta cell loss and hyperglycaemia. *Diabetologia* 50: 142–150, 2007.
14. Dallas-Yang Q, Shen X, Strowski M, Brady E, Saperstein R, Gibson RE, Szalkowski D, Qureshi SA, Candelore MR, Fenyk-Melody JE, Parmee ER, Zhang BB, Jiang G. Hepatic glucagon receptor binding and glucose-lowering in vivo by peptidyl and non-peptidyl glucagon receptor antagonists. *Eur J Pharmacol* 501: 225–234, 2004.
15. de Heer J, Rasmussen C, Coy DH, Holst JJ. Glucagon-like peptide-1, but not glucose-dependent insulinotropic peptide, inhibits glucagon secretion via somatostatin (receptor subtype 2) in the perfused rat pancreas. *Diabetologia* 51: 2263–2270, 2008.
16. Dinneen S, Alzaid A, Turk D, Rizza R. Failure of glucagon suppression contributes to postprandial hyperglycaemia in IDDM. *Diabetologia* 38: 337–343, 1995.
17. Drucker DJ, Asa S. Glucagon gene expression in vertebrate brain. *J Biol Chem* 263: 13475–13478, 1988.
18. Dunning BE, Gerich JE. The role of alpha-cell dysregulation in fasting and postprandial hyperglycemia in type 2 diabetes and therapeutic implications. *Endocr Rev* 28: 253–283, 2007.
19. Gastaldelli A, Baldi S, Pettiti M, Toschi E, Camastra S, Natali A, Landau BR, Ferrannini E. Influence of obesity and type 2 diabetes on gluconeogenesis and glucose output in humans: a quantitative study. *Diabetes* 49: 1367–1373, 2000.
20. Geary N, Le Sauter J, Noh U. Glucagon acts in the liver to control spontaneous meal size in rats. *Am J Physiol Regul Integr Comp Physiol* 264: R116–R122, 1993.
21. Gelling RW, Du XQ, Dichmann DS, Romer J, Huang H, Cui L, Obici S, Tang B, Holst JJ, Fledelius C, Johansen PB, Rossetti L, Jelicks LA, Serup P, Nishimura E, Charron MJ. Lower blood glucose, hyperglucagonemia, and pancreatic alpha cell hyperplasia in glucagon receptor knockout mice. *Proc Natl Acad Sci USA* 100: 1438–1443, 2003.
22. Gerich JE, Lorenzi M, Karam JH, Schneider V, Forsham PH. Abnormal pancreatic glucagon secretion and postprandial hyperglycemia in diabetes mellitus. *JAMA* 234: 159–165, 1975.

23. Gonzalez-Munoz C, Nieto-Ceron S, Cabezas-Herrera J, Hernandez-Cascales J. Glucagon increases contractility in ventricle but not in atrium of the rat heart. *Eur J Pharmacol* 587: 243–247, 2008.
24. Gravholt CH, Moller N, Jensen MD, Christiansen JS, Schmitz O. Physiological levels of glucagon do not influence lipolysis in abdominal adipose tissue as assessed by microdialysis. *J Clin Endocrinol Metab* 86: 2085–2089, 2001.
25. Guettet C, Rostaqui N, Mathe D, Lecuyer B, Navarro N, Jacotot B. Effect of chronic glucagon administration on lipoprotein composition in normally fed, fasted and cholesterol-fed rats. *Lipids* 26: 451–458, 1991.
26. Gustavson SM, Chu CA, Nishizawa M, Neal D, Farmer B, Yang Y, Donahue EP, Flakoll P, Cherrington AD. Effects of hyperglycemia, glucagon, and epinephrine on renal glucose release in the conscious dog. *Metabolism* 53: 933–941, 2004.
27. Hansen LH, Abrahamsen N, Nishimura E. Glucagon receptor mRNA distribution in rat tissues. *Peptides* 16: 1163–1166, 1995.
28. Harney JA, Rodgers RL. Insulin-like stimulation of cardiac fuel metabolism by physiological levels of glucagon: involvement of PI3K but not cAMP. *Am J Physiol Endocrinol Metab* 295: E155–E161, 2008.
29. Heckemeyer CM, Barker J, Duckworth WC, Solomon SS. Studies of the biological effect and degradation of glucagon in the rat perfused isolated adipose cell. *Endocrinology* 113: 270–276, 1983.
30. Henkel E, Menschikowski M, Koehler C, Leonhardt W, Hanefeld M. Impact of glucagon response on postprandial hyperglycemia in men with impaired glucose tolerance and type 2 diabetes mellitus. *Metabolism* 54: 1168–1173, 2005.
31. Hoosain NM, Gurd RS. Identification of glucagon receptors in rat brain. *Proc Natl Acad Sci USA* 81: 4368–4372, 1984.
32. Inokuchi A, Oomura Y, Nishimura H. Effect of intracerebroventricularly infused glucagon on feeding behavior. *Physiol Behav* 33: 397–400, 1984.
33. Jelinek LJ, Lok S, Rosenberg GB, Smith RA, Grant FJ, Biggs S, Bensch PA, Kuijper JL, Sheppard PO, Sprecher CA, O'Hara PJ, Foster D, Walker KM, Chen LH, Mckernan PA, Kindsvogel W. Expression cloning and signaling properties of the rat glucagon receptor. *Science* 259: 1614–1616, 1993.
34. Jiang G, Zhang BB. Glucagon and regulation of glucose metabolism. *Am J Physiol Endocrinol Metab* 284: E671–E678, 2003.
35. Jin SL, Han VK, Simmons JG, Towle AC, Lauder JM, Lund PK. Distribution of glucagonlike peptide I (GLP-I), glucagon, and glicentin in the rat brain: an immunocytochemical study. *J Comp Neurol* 271: 519–532, 1988.
36. Johnson DG, Goebel CU, Hruby VJ, Bregman MD, Trivedi D. Hyperglycemia of diabetic rats decreased by a glucagon receptor antagonist. *Science* 215: 1115–1116, 1982.
37. Kieffer TJ, Heller RS, Unson CG, Weir GC, Habener JF. Distribution of glucagon receptors on hormone-specific endocrine cells of rat pancreatic islets. *Endocrinology* 137: 5119–5125, 1996.
38. Kim RM, Chang J, Lins AR, Brady E, Candelore MR, Dallas-Yang Q, Ding V, Dragovic J, Iliff S, Jiang G, Mock S, Qureshi S, Saperstein R, Szalkowski D, Tamvakopoulos C, Tota L, Wright M, Yang X, Tata JR, Chapman K, Zhang BB, Parmee ER. Discovery of potent, orally active benzimidazole glucagon receptor antagonists. *Bioorg Med Chem Lett* 18: 3701–3705, 2008.
39. Lau J, Behrens C, Sidelmann UG, Knudsen LB, Lundt B, Sams C, Ynddal L, Brand CL, Pridal L, Ling A, Kiel D, Plewe M, Shi S, Madsen P. New beta-alanine derivatives are orally available glucagon receptor antagonists. *J Med Chem* 50: 113–128, 2007.
40. Li XC, Liao TD, Zhuo JL. Long-term hyperglucagonaemia induces early metabolic and renal phenotypes of type 2 diabetes in mice. *Clin Sci (Lond)* 114: 591–601, 2008.
41. Li Y, Hansotia T, Yusta B, Ris F, Halban PA, Drucker DJ. Glucagon-like peptide-1 receptor signaling modulates beta cell apoptosis. *J Biol Chem* 278: 471–478, 2003.
42. Liang R, Abrardo L, Brady EJ, Candelore MR, Ding V, Saperstein R, Tota LM, Wright M, Mock S, Tamvakopolous C, Tong S, Zheng S, Zhang BB, Tata JR, Parmee ER. Design and synthesis of conformationally constrained tri-substituted ureas as potent antagonists of the human glucagon receptor. *Bioorg Med Chem Lett* 17: 587–592, 2007.
43. Liang Y, Osborne MC, Monia BP, Bhanot S, Gaarde WA, Reed C, She P, Jetton TL, Demarest KT. Reduction in glucagon receptor expression by an antisense oligonucleotide ameliorates diabetic syndrome in db/db mice. *Diabetes* 53: 410–417, 2004.
44. Lins PE, Wajngot A, Adamson U, Vranic M, Efendic S. Minimal increases in glucagon levels enhance glucose production in man with partial hypoinsulinemia. *Diabetes* 32: 633–636, 1983.
45. Liu Y, Dentin R, Chen D, Hedrick S, Ravnskjaer K, Schenk S, Milne J, Meyers DJ, Cole P, Yates J 3rd, Olefsky J, Guarente L, Montminy M. A fasting inducible switch modulates gluconeogenesis via activator/coactivator exchange. *Nature* 456: 269–273, 2008.
46. Longuet C, Sinclair EM, Maida A, Baggio LL, Maziarz M, Charron MJ, Drucker DJ. The Glucagon receptor is required for the adaptive metabolic response to fasting. *Cell Metab* 8: 359–371, 2008.
47. Ma X, Zhang Y, Gromada J, Sewing S, Berggren PO, Buschard K, Salehi A, Vikman J, Rorsman P, Eliasson L. Glucagon stimulates exocytosis in mouse and rat pancreatic alpha-cells by binding to glucagon receptors. *Mol Endocrinol* 19: 198–212, 2005.
48. MacDonald PE, De Marinis YZ, Ramrachea R, Salehi A, Ma X, Johnson PR, Cox R, Eliasson L, Rorsman P. A K ATP channel-dependent pathway within alpha cells regulates glucagon release from both rodent and human islets of Langerhans. *PLoS Biol* 5: e143, 2007.
49. Madsen P, Ling A, Plewe M, Sams CK, Knudsen LB, Sidelmann UG, Ynddal L, Brand CL, Andersen B, Murphy D, Teng M, Truesdale L, Kiel D, May J, Kuki A, Shi S, Johnson MD, Teston KA, Feng J, Lakis J, Anderes K, Gregor V, Lau J. Optimization of alkylidene hydrazide based human glucagon receptor antagonists. Discovery of the highly potent and orally available 3-cyano-4-hydroxybenzoic acid [1-(2,3,5,6-tetramethylbenzyl)-1H-indol-4-ylmethylene]hydrazide. *J Med Chem* 45: 5755–5775, 2002.
50. Marubashi S, Tominaga M, Katagiri T, Yamatani K, Yawata Y, Hara M, Sasaki H. Hyperglycemic effect of glucagon administered intracerebroventricularly in the rat. *Acta Endocrinol (Copenh)* 108: 6–10, 1985.
51. Miki T, Liss B, Minami K, Shiuchi T, Saraya A, Kashima Y, Horiuchi M, Ashcroft F, Minokoshi Y, Roeper J, Seino S. ATP-sensitive K⁺ channels in the hypothalamus are essential for the maintenance of glucose homeostasis. *Nat Neurosci* 4: 507–512, 2001.
52. Moens K, Flamez D, Van Schravendijk C, Ling Z, Pipeleers D, Schuit F. Dual glucagon recognition by pancreatic beta-cells via glucagon and glucagon-like peptide 1 receptors. *Diabetes* 47: 66–72, 1998.
53. Moens K, Heimberg H, Flamez D, Huybens P, Quartier E, Ling Z, Pipeleers D, Gremlich S, Thorens B, Schuit F. Expression and functional activity of glucagon, glucagon-like peptide 1 and glucose-dependent insulinotropic peptide receptors in rat pancreatic islet cells. *Diabetes* 45: 257–261, 1996.
54. Mulvehill JB, Hui YS, Barnes LD, Palumbo PJ, Dousa TP. Glucagon-sensitive adenylate cyclase in human renal medulla. *J Clin Endocrinol Metab* 42: 380–384, 1976.
55. Nafikov RA, Ametaj BN, Bohe G, Koehler KJ, Young JW, Beitz DC. Prevention of fatty liver in transition dairy cows by subcutaneous injections of glucagon. *J Dairy Sci* 89: 1533–1545, 2006.
56. Parker JC, Andrews KM, Allen MR, Stock JL, McNeish JD. Glycemic control in mice with targeted disruption of the glucagon receptor gene. *Biochem Biophys Res Commun* 290: 839–843, 2002.
57. Parnaud G, Bosco D, Berney T, Pattou F, Kerr-Conte J, Donath MY, Bruun C, Mandrup-Poulsen T, Billestrup N, Halban PA. Proliferation of sorted human and rat beta cells. *Diabetologia* 51: 91–100, 2008.
58. Patel GK, Whalen GE, Soergel KH, Wu WC, Meade RC. Glucagon effects on the human small intestine. *Dig Dis Sci* 24: 501–508, 1979.
59. Perea A, Clemente F, Martinell J, Villanueva-Penacarrillo ML, Valverde I. Physiological effect of glucagon in human isolated adipocytes. *Horm Metab Res* 27: 372–375, 1995.
60. Petersen KF, Sullivan JT. Effects of a novel glucagon receptor antagonist (Bay 27-9955) on glucagon-stimulated glucose production in humans. *Diabetologia* 44: 2018–2024, 2001.
61. Quesada I, Tuduri E, Ripoll C, Nadal A. Physiology of the pancreatic alpha-cell and glucagon secretion: role in glucose homeostasis and diabetes. *J Endocrinol* 199: 5–19, 2008.
62. Shah P, Basu A, Basu R, Rizza R. Impact of lack of suppression of glucagon on glucose tolerance in humans. *Am J Physiol Endocrinol Metab* 277: E283–E290, 1999.
63. Shah P, Vella A, Basu A, Basu R, Schwenk WF, Rizza RA. Lack of suppression of glucagon contributes to postprandial hyperglycemia in subjects with type 2 diabetes mellitus. *J Clin Endocrinol Metab* 85: 4053–4059, 2000.
64. Sinclair EM, Yusta B, Streutker C, Baggio LL, Koehler J, Charron MJ, Drucker DJ. Glucagon receptor signaling is essential for control of murine hepatocyte survival. *Gastroenterology* 135: 2096–2106, 2008.

65. **Singh V, Grotzinger C, Nowak KW, Zacharias S, Goncz E, Pless G, Sauer IM, Eichhorn I, Pfeiffer-Guglielmi B, Hamprecht B, Wiedemann B, Plockinger U, Strowski MZ.** Somatostatin receptor subtype-2-deficient mice with diet-induced obesity have hyperglycemia, nonfasting hyperglucagonemia, and decreased hepatic glycogen deposition. *Endocrinology* 148: 3887–3899, 2007.
66. **Sloop KW, Cao JX, Siesky AM, Zhang HY, Bodenmiller DM, Cox AL, Jacobs SJ, Moyers JS, Owens RA, Showalter AD, Brenner MB, Raap A, Gromada J, Berridge BR, Monteith DK, Porksen N, McKay RA, Monia BP, Bhanot S, Watts LM, Michael MD.** Hepatic and glucagon-like peptide-1-mediated reversal of diabetes by glucagon receptor antisense oligonucleotide inhibitors. *J Clin Invest* 113: 1571–1581, 2004.
67. **Sorensen H, Brand CL, Neschen S, Holst JJ, Fosgerau K, Nishimura E, Shulman GI.** Immunoneutralization of endogenous glucagon reduces hepatic glucose output and improves long-term glycemic control in diabetic ob/ob mice. *Diabetes* 55: 2843–2848, 2006.
68. **Sorensen H, Winzell MS, Brand CL, Fosgerau K, Gelling RW, Nishimura E, Ahren B.** Glucagon receptor knockout mice display increased insulin sensitivity and impaired beta-cell function. *Diabetes* 55: 3463–3469, 2006.
69. **Taylor I, Duthie HL, Cumberland DC, Smallwood R.** Glucagon and the colon. *Gut* 16: 973–978, 1975.
70. **Unson CG, Gurzenda EM, Merrifield RB.** Biological activities of des-His1[Glu9]glucagon amide, a glucagon antagonist. *Peptides* 10: 1171–1177, 1989.
71. **Vuguin PM, Kedees MH, Cui L, Guz Y, Gelling RW, Nejathaim M, Charron MJ, Teitelman G.** Ablation of the glucagon receptor gene increases fetal lethality and produces alterations in islet development and maturation. *Endocrinology* 147: 3995–4006, 2006.
72. **Weatherford SC, Ritter S.** Lesion of vagal afferent terminals impairs glucagon-induced suppression of food intake. *Physiol Behav* 43: 645–650, 1988.
73. **White CM.** A review of potential cardiovascular uses of intravenous glucagon administration. *J Clin Pharmacol* 39: 442–447, 1999.
74. **Xu E, Kumar M, Zhang Y, Ju W, Obata T, Zhang N, Liu S, Wendt A, Deng S, Ebina Y, Wheeler MB, Braun M, Wang Q.** Intra-islet insulin suppresses glucagon release via GABA-GABA(A) receptor system. *Cell Metab* 3: 47–58, 2006.
75. **Yano Y, Cesar KR, Araujo M, Rodrigues AC Jr, Andrade LC, Magaldi AJ.** Aquaporin 2 expression increased by glucagon in normal rat inner medullary collecting ducts. *Am J Physiol Renal Physiol* 296: F54–F59, 2009.
76. **Zhou H, Zhang T, Harmon JS, Bryan J, Robertson RP.** Zinc, not insulin, regulates the rat alpha-cell response to hypoglycemia in vivo. *Diabetes* 56: 1107–1112, 2007.

