

Altered cAMP and Ca²⁺ Signaling in Mouse Pancreatic Islets With Glucagon-Like Peptide-1 Receptor Null Phenotype

Daisy Flamez, Patrick Gilon, Karen Moens, An Van Breusegem, Dominique Delmeire, Louise A. Scrocchi, Jean-Claude Henquin, Daniel J. Drucker, and Frans Schuit

β-Cells from rodents and humans express different receptors recognizing hormones of the secretin-glucagon family, which—when activated—synergize with glucose in the control of insulin release. We have recently reported that isolated islets from mice homozygous for a GLP-1 receptor null mutation (GLP-1R^{-/-}) exhibit a well-preserved insulin-secretory response to glucose. This observation can be interpreted in two different ways: 1) the presence of GLP-1R is not essential for the secretory response of isolated islets to glucose alone; 2) β-cells in GLP-1R^{-/-} pancreases underwent compensatory changes in response to the null mutation. To explore these possibilities, we studied islets from control GLP-1R^{+/+} mice in the absence or presence of 1 μmol/l exendin (9–39)amide, a specific and potent GLP-1R antagonist. Exendin (9–39)amide (15-min exposure) reduced glucose-induced insulin secretion from both perfused and statically incubated GLP-1R^{+/+} islets by 50% ($P < 0.05$), and reduced islet cAMP production in parallel ($P < 0.001$). Furthermore, GLP-1R^{-/-} islets exhibited: 1) reduced cAMP accumulation in the presence of 20 mmol/l glucose (knockout islets versus control islets, 12 ± 1 vs. 27 ± 3 fmol · islet⁻¹ · 15 min⁻¹; $P < 0.001$) and exaggerated acceleration of cAMP production by 10 nmol/l glucose-dependent insulinotropic peptide (GIP) (increase over 20 mmol/l glucose by GIP in knockout islets versus control islets: 66 ± 5 vs. 14 ± 3 fmol · islet⁻¹ · 15 min⁻¹; $P < 0.001$); 2) increased mean cytosolic [Ca²⁺]_c ([Ca²⁺]_c) at 7, 10, and 15 mmol/l glucose in knockout islets versus control islets; and 3) signs of asynchrony of [Ca²⁺]_c oscillations between different islet subregions. In conclusion, disruption of GLP-1R signaling is associated with reduced basal but enhanced GIP-stimulated cAMP production and abnormalities in basal and glucose-stimulated [Ca²⁺]_c. These abnormali-

ties suggest that GLP-1R signaling is an essential upstream component of multiple β-cell signaling pathways. *Diabetes* 48:1979–1986, 1999

Oral glucose administration stimulates insulin secretion more than intravenous glucose infusion (1–3) does. This so-called incretin effect of oral glucose results from the secretion of gut hormones that potentiate the effect of glucose on insulin secretion (4). Two intestinal peptides released during a meal are important in mediating the incretin effect: GIP (glucose-dependent insulinotropic peptide) and GLP-1 [glucagon-like peptide-1-(7–36) amide] (4,5). Evidence for the functional importance of GIP and GLP-1 as physiologically relevant incretins derives from a combination of experimental approaches including immunoneutralization of GIP action (6), blockade of GLP-1 action with the specific antagonist exendin (9–39)amide (7), and studies of GLP-1 receptor (GLP-1R) ^{-/-} mice (8). Because both peptides exert their potentiating effects on insulin release only at elevated glucose concentrations, they have been considered as potential therapeutic agents for patients with type 2 diabetes (4,9).

It has also been suggested that regular exposure of β-cells to GLP-1 is important to maintain normal insulin secretory responsiveness to glucose (10,11). However, we have recently shown that isolated islets from mice homozygous for a GLP-1 receptor null mutation (GLP-1R^{-/-}) exhibit a well-preserved insulin response to glucose (12). These observations suggest that either the presence of GLP-1R is not required for glucose competence in isolated islets or that mice with GLP-1R^{-/-} have compensated for their genetic defect. One element of compensation appears to involve the GIP component of the enteroinsular axis, inasmuch as both GIP secretion and GIP action are increased in GLP-1R^{-/-} mice (13).

In this study we have explored the possibility that additional compensatory changes occur at the level of GLP-1R^{-/-} islets following disruption of GLP-1 signaling. Previous *in vitro* experiments showed that both GLP-1 and GIP stimulate insulin secretion mainly by increasing cellular cAMP levels in β-cells (14,15) or β-cell lines (16,17) but other signaling pathways may also be involved (18). Compensation might therefore occur at the level of intracellular cAMP, which is critical for insulin release (19–22) as well as cytosolic [Ca²⁺]_c ([Ca²⁺]_c), which is generally accepted to be one of the main regulators of exocytosis (23,24). Our data

From the Diabetes Research Center (D.F., K.M., A.V.B., D.D., F.S.), Vrije Universiteit Brussel; the Unité d'Endocrinologie et Métabolisme (P.G., J.-C.H.), Université Catholique de Louvain, Brussels, Belgium; and the Department of Medicine (L.A.S., D.J.D.), University of Toronto, and Banting and Best Diabetes Center, The Toronto Hospital, Toronto, Ontario, Canada.

Address correspondence and reprint requests to Dr. Frans Schuit, Department of Biochemistry, Diabetes Research Center, Faculty of Medicine, Vrije Universiteit Brussel, Laarbeeklaan 103, B-1090 Brussels, Belgium. E-mail: fschuit@minf.vub.ac.be.

Received for publication 8 December 1998 and accepted in revised form 17 June 1999.

D.J.D. has been a consultant for Amylin Pharmaceuticals.

Authors D.F. and P.G. contributed equally to this study.

ACh, acetylcholine; BSA, bovine serum albumin; [Ca²⁺]_c, cytosolic [Ca²⁺]; GIP, glucose-dependent insulinotropic peptide; GLP-1, glucagon-like peptide-1-(7–36) amide; GLP-1R, GLP-1 receptor; IBMX, 3-isobutyl-1-methylxanthine; Sp-cAMPS, cAMP Sp-isomer.




FIG. 1. Effect of exendin (9-39)amide on glucose-induced insulin release. Islets of female GLP-1R^{+/+} mice were perfused with 10 mmol/l glucose, 20 mmol/l glucose, and 20 mmol/l glucose with 10 nmol/l GLP-1. One batch of islets was perfused with 1 μ mol/l exendin (9-39)amide (●), the other without exendin (9-39)amide (□). Data represent the means \pm SE of four experiments. For graphical clarity, mean symbols and error bars are shown once per four consecutive minute fractions.

indicate that changes in both cAMP and [Ca²⁺]_i occur in the GLP-1R^{-/-} model.

RESEARCH DESIGN AND METHODS

Chemicals. Synthetic GLP-1 (7-36)amide and GIP were purchased from Sigma (St. Louis, MO); 3-isobutyl-1-methylxanthine (IBMX) was purchased from Aldrich (Janssen Chimica, Beerse, Belgium). Exendin (9-39)amide was a gift from Dr. J. Vandekerckhove (Department of Physiological Chemistry, University of Ghent, Ghent, Belgium). cAMP Sp-isomer (Sp-cAMPS) was purchased from Biolog Life Science Institute (Bremen, Germany). All other products were of analytical grade and obtained from either Merck (Darmstadt, Germany) or Sigma.

Animals. The study was conducted on 8- to 12-week-old male and female CD1 mice fed ad libitum that were wild type (GLP-1R^{+/+}) or homozygous (GLP-1R^{-/-}) for a targeted null mutation of the *GLP-1R* gene (8). The animals were bred under specific pathogen-free conditions and cared for according to Belgian regulations of animal welfare.

Isolation and culture of mouse pancreatic islets. Islets of Langerhans were isolated as described previously (12) using collagenase P (Boehringer Mannheim, Mannheim, Germany). Isolated islets were then cultured overnight in F10 medium Nutrient Mixture (Ham) (GIBCO BRL, Life Technologies, Strathclyde, U.K.) containing 0.075 mg/ml penicillin, 0.1 mg/ml streptomycin, 0.5% (wt/vol) bovine serum albumin (BSA) (fraction V, radioimmunoassay grade; Sigma), 2 mmol/l glutamine, and 10 mmol/l glucose (25). Insulin release and islet cAMP experiments were performed after overnight culture, and the measurements of [Ca²⁺]_i were performed after two days of culture. Experiments using control islets which were cultured for one or two days gave similar results.

Insulin release. Perfusion experiments were performed at 37°C with overnight cultured female GLP-1R^{+/+} control islets using a perfusion system with a multiple microchamber module (Endotronics, Coon Rapids, MN) (12). Islets were divided in equal batches (~200 islets per batch) and perfused in parallel, with various glucose concentrations with or without 10 nmol/l GLP-1. One perfusion column was exposed during the entire experiment to 1 μ mol/l exendin (9-39)amide. Preperfusion (15 min) was done in basal medium that consisted of F10 medium Nutrient Mixture supplemented with 0.5% BSA, 2 mmol/l glutamine, 2 mmol/l CaCl₂, and 1 mmol/l glucose equilibrated with 95% O₂/5% CO₂ at a flow rate of 0.5 ml/min. Samples were collected every minute and assayed for immunoreactive insulin (19). The dead space was taken into account when expressing results as a function of time. Insulin content of the GLP-1R^{+/+} islets (63 \pm 8 ng/islet) (*n* = 10) was measured after perfusion by sonicating the Biogel P2 containing the islets in 5 ml of 2 mmol/l acetic acid/0.25% BSA.




FIG. 2. cAMP content measured in the presence of 250 μ mol/l IBMX (A) and insulin release measured without IBMX (B) from GLP-1R^{+/+} and GLP-1R^{-/-} islets. The effects of 20 mmol/l glucose alone or 20 mmol/l glucose plus exendin (9-39)amide (1 μ mol/l), GLP-1 (10 nmol/l), GIP (10 nmol/l), and GIP plus exendin (9-39)amide were assessed as indicated using parallel incubations of GLP-1R^{+/+} (□) and GLP-1R^{-/-} islets (■). Data are means \pm SE of 13 (GLP-1R^{+/+} islets) and 7 (GLP-1R^{-/-} islets) independent experiments.

The effect of 1 μ mol/l exendin (9-39)amide was also studied on 20 mmol/l glucose-induced insulin release from control islets during 15 min static incubations. Furthermore, effects of exendin (9-39)amide, GLP-1, GIP, and Sp-cAMPS were tested during 60-min static incubations in the presence of 20 mmol/l glucose using triplicate batches of 20 islets isolated from GLP-1R^{+/+} and GLP-1R^{-/-} mice. For this purpose, overnight cultured islets were washed three times in Earle's HEPES medium (124 mmol/l NaCl, 5.4 mmol/l KCl, 1.8 mmol/l CaCl₂, 0.8 mmol/l MgSO₄, 1 mmol/l NaH₂PO₄, 14 mmol/l NaHCO₃, 10 mmol/l HEPES, and 0.5% BSA) (no glucose). Incubation was started by adding fresh Earle's HEPES medium supplemented with test agents at final concentrations that are indicated in Figs. 2B and 3. Results were expressed as percentage of the insulin content, measured in each individual batch of islets after finishing the experiment. No difference in insulin content of GLP-1R^{+/+} and ^{-/-} islets was observed after the static incubations: 67 \pm 4 vs. 65 \pm 3 ng/islet (mean \pm SE of 13 experiments).

Measurement of cAMP production. Levels of cAMP were determined in whole islets as was described previously (15,20). Batches of 40 islets were incubated for 15 min at 37°C in Earle's HEPES medium supplemented with 0.5% BSA, 250 μ mol/l IBMX, 20 mmol/l glucose, and the peptide hormones as indicated. cAMP content of acetylated samples was measured using a commercially available [¹²⁵I]-cAMP radioimmunoassay kit (Amersham, Little Chalfont, U.K.).

Measurements of [Ca²⁺]_i. A bicarbonate-buffered medium (pH 7.4) containing 120 mmol/l NaCl, 4.8 mmol/l KCl, 2.5 mmol/l CaCl₂, 1.2 mmol/l MgCl₂, and 24 mmol/l NaHCO₃, supplemented with 1 mg/ml BSA (fraction V; Boehringer Mannheim), gassed with 94% O₂/6% CO₂, was used for Ca²⁺ experiments. In one experimental condition where the KCl concentration was increased to 30 mmol/l, that of NaCl was decreased accordingly to keep the osmolarity constant. After two days of culture, male islets were loaded with 2 μ mol/l fura-PE3/AM (Teflabs, Austin, TX) for 90-120 min at 37°C in a bicarbonate-buffered solution containing 10 mmol/l glucose and transferred into a temperature-controlled perfusion chamber (37°C) with a bottom made of a glass coverslip. The flow rate of perfusion was 1.3 ml/min. The islets were excited successively at 340 and 380 nm, the fluorescence emitted at 510 nm was captured by a CCD camera (Photonic Science, Tunbridge Wells, U.K.), and the images were analyzed by the QuantCell system (Applied Imaging, Sunderland, U.K.). The technique has previously been described in detail (26). In the experiments illustrated in Fig. 6, the 510-nm




FIG. 3. Effect of the cAMP analog Sp-cAMPS on glucose-induced insulin release in GLP-1R^{+/+} and GLP-1R^{-/-} islets. Batches of 20 islets isolated from GLP-1R^{+/+} (□) and GLP-1R^{-/-} mice (■) were challenged with different concentrations of Sp-cAMPS as indicated in the presence of 20 mmol/l glucose. Data are means \pm SE of five independent experiments.

fluorescence was monitored by a photomultiplier-based system (Photon Technology International, Princeton, NJ).

Statistical analysis. The results are illustrated either as means \pm SE of at least three different experiments or as representative data obtained with the indicated number of islets. The statistical significance of observed differences was assessed by unpaired two-tailed Student's *t* tests.

RESULTS

Effect of exendin (9–39)amide on glucose-induced insulin secretion. Because glucose-induced insulin release is well preserved in GLP-1R^{-/-} islets (12), we investigated the hypothesis that the GLP-1R is not essential for a normal glucose-induced insulin secretion by assessing the effect of pharmacological inhibition of the GLP-1R with 1 μ mol/l exendin (9–39)amide in wild-type islets. As is shown in Fig. 1, this treatment almost completely prevented the potentiation by 10 nmol/l GLP-1 (7–36)amide of 20 mmol/l glucose-induced insulin release [3.5 ± 1.5 -fold potentiation in controls vs. 1.5 ± 0.2 -fold potentiation in the presence of 1 μ mol/l exendin (9–39)amide; $P < 0.05$]. Furthermore, glucose-induced insulin release was reduced by 40–50% in exendin (9–39)amide-treated islets: release above basal per 15 min for control and exendin (9–39)amide treatment was 0.34 ± 0.04 vs. $0.14 \pm 0.04\%$ (10 mmol/l glucose) and 0.53 ± 0.06 vs. $0.31 \pm 0.07\%$ (20 mmol/l glucose) of islet insulin content, respectively ($P < 0.05$). A similar reduction in 20 mmol/l glucose-induced insulin release was observed in statically incubated control islets treated for 15 min with exendin (9–39)amide (glucose alone: $1.7 \pm 0.2\%$ vs. glucose plus exendin (9–39)amide: $0.8 \pm 0.2\%$ of islet insulin content; mean \pm SE, $n = 3$, $P < 0.05$). These observations implicate a role for the GLP-1R in the response of β -cells to glucose stimulation in vitro, and suggest that the preserved glucose-induced insulin secretion in GLP-1R^{-/-} islets is due to adaptive mechanisms that compensate for the loss of GLP-1R activity in β -cells.

Islet cAMP accumulation and insulin release from islets isolated from GLP-1R^{+/+} and GLP-1R^{-/-} mice.

Islet cAMP accumulation, measured in the presence of 20 mmol/l glucose and 250 μ mol/l IBMX (Fig. 2A), was two times higher in GLP-1R^{+/+} islets (27 ± 3 fmol \cdot islet⁻¹ \cdot 15 min⁻¹) than in GLP-1R^{-/-} islets (12 ± 1 fmol \cdot islet⁻¹ \cdot 15 min⁻¹; $P < 0.01$). Exendin (9–39)amide did not affect islet cAMP accumulation in GLP-1R^{-/-} islets, but it decreased cAMP accumulation in the GLP-1R^{+/+} islets (12 ± 1 fmol \cdot islet⁻¹ \cdot 15




FIG. 4. Effect of glucose, ACh, diazoxide, and high potassium on $[Ca^{2+}]_i$ in GLP-1R^{+/+} and GLP-1R^{-/-} islets. The glucose concentration of the perfusion medium was increased from 3 (G3) to 15 mmol/l (G15) after 1 min of recording. ACh (1 μ mol/l), diazoxide (250 μ mol/l), and 30 mmol/l K⁺ (K30) were applied as indicated. The traces are representative or mean results from 38 GLP-1R^{+/+} and 38 GLP-1R^{-/-} islets.

min⁻¹; $P < 0.001$), so that in the presence of exendin (9–39)amide, cAMP levels in GLP-1R^{-/-} and GLP-1R^{+/+} islets were comparable. These findings implicate a role for GLP-1R signaling in the regulation of basal islet cAMP levels in vitro. Paradoxically, exendin (9–39)amide did not affect glucose-induced insulin secretion in GLP-1R^{+/+} islets after 15 min incubation in the presence of IBMX [$3.1 \pm 0.5\%$ of insulin content in exendin (9–39)amide-treated islets vs. $3.7 \pm 0.6\%$ in control islets, $n = 4$] or after 60 min incubation in the absence of IBMX (Fig. 2B). After 15-min incubations without IBMX, however, 20 mmol/l glucose-induced insulin secretion was inhibited by exendin (9–39)amide ($0.8 \pm 0.2\%$ of insulin content vs. $1.7 \pm 0.2\%$ of insulin content in glucose alone, $n = 4$, $P < 0.05$). This short-term inhibition is consistent with the results obtained with the perfused islets (Fig. 1).

GLP-1 slightly stimulated ($P = 0.01$) cAMP accumulation in control islets incubated in the presence of 20 mmol/l glucose and 250 μ mol/l IBMX (Fig. 2A). The accompanying increase in insulin secretion, however, failed to reach statistical significance (5.1 ± 0.7 vs. $3.7 \pm 0.6\%$ of islet content, $n = 4$, $P = 0.12$). A larger effect of GLP-1 on insulin secretion was observed in islets incubated for 60 min in absence of IBMX (Fig. 2B).

It was previously reported that GLP-1R^{-/-} islets exhibit an increased responsiveness to GIP (13). This behavior is again shown in Fig. 2B, where GLP-1R^{-/-} islets respond more to 20 mmol/l glucose plus 10 nmol/l GIP than do control islets (1.5-fold increase above basal in the control islets vs. threefold stimulation in GLP-1R^{-/-} islets, $P < 0.001$). This difference in insulin secretion between GLP-1R^{-/-} and GLP-1R^{+/+} islets is paralleled by a larger GIP-induced formation of cAMP (1.5-fold increase of basal in control islets [$P = 0.001$] vs. 6.4-fold stimulation above basal in GLP-1R^{-/-} islets, $P = 0.0001$). In the presence of 1 $\mu\text{mol/l}$ exendin (9–39)amide, cAMP production in GIP-treated control mouse islets was slightly lower ($P < 0.01$) than in islets treated with GIP but with no exendin (9–39)amide. This difference can be explained, however, by the effect of 1 $\mu\text{mol/l}$ exendin (9–39)amide on 20 mmol/l glucose-induced cAMP production. Thus, the GIP-induced increase in cAMP production above that with 20 mmol/l glucose alone was $14 \pm 3 \text{ fmol} \cdot \text{islet}^{-1} \cdot 15 \text{ min}^{-1}$ without exendin (9–39)amide vs. $21 \pm 2 \text{ fmol} \cdot \text{islet}^{-1} \cdot 15 \text{ min}^{-1}$ in the presence of exendin (9–39)amide. Second, the GIP-induced cAMP accumulation in knockout islets was also unaffected by exendin (9–39)amide, the increase of GIP above 20 mmol/l glucose alone being $66 \pm 5 \text{ fmol} \cdot \text{islet}^{-1} \cdot 15 \text{ min}^{-1}$ without exendin (9–39)amide vs. $59 \pm 3 \text{ fmol} \cdot \text{islet}^{-1} \cdot 15 \text{ min}^{-1}$ with exendin (9–39)amide. Third, the increase in insulin secretion induced by GIP in GLP-1R^{-/-} islets was similar in the presence and absence of exendin (9–39)amide. In summary, 1 $\mu\text{mol/l}$ exendin (9–39)amide does not interfere with GIP signaling in mouse islets.

Effect of the cAMP analog Sp-cAMPS on 20 mmol/l glucose-induced insulin release. One possible explanation for the similar secretory response of GLP-1R^{-/-} and GLP-1R^{+/+} islets in the presence of a twofold difference in islet cAMP content is that the β -cells from GLP-1R^{-/-} mice have augmented the sensitivity of their secretory machinery to cAMP. To investigate this possibility, we exposed batches of islets to different concentrations of the cAMP analog Sp-cAMPS, a potent stimulator of protein kinase A (Fig. 3). Insulin secretion at 20 mmol/l glucose was again comparable in GLP-1R^{+/+} and GLP-1R^{-/-} islets. Furthermore, Sp-cAMPS produced a concentration-dependent potentiation of insulin secretion without any suggestion of an increased sensitivity of the insulin secretory responsiveness to cAMP in the GLP-1R^{-/-} islets.

Effects of glucose, acetylcholine, and high K⁺ on islet [Ca²⁺]_i. Because [Ca²⁺]_i and cAMP synergize at the level of exocytosis (27), we assessed the possibility that a larger [Ca²⁺]_i response could compensate for the lower levels of cAMP, resulting in well-preserved glucose-induced insulin secretion in GLP-1R^{-/-} islets. In 3 mmol/l glucose, [Ca²⁺]_i was low, stable, and similar in control and GLP-1R^{-/-} islets (Fig. 4). Raising the glucose concentration to 15 mmol/l produced a biphasic increase in [Ca²⁺]_i. The kinetics and amplitude of the initial [Ca²⁺]_i phase were similar in control and GLP-1R^{-/-} islets. In contrast, the characteristics of the second phase were different in the two types of islets. In control islets, [Ca²⁺]_i displayed large amplitude and regular oscillations above a slightly elevated level (Fig. 4A). In GLP-1R^{-/-} islets, [Ca²⁺]_i oscillations occurred from a higher level, but were of smaller amplitude and poorly organized (Fig. 4B). The overall [Ca²⁺]_i response to glucose was significantly larger in GLP-1R^{-/-} ($215 \pm 4 \text{ nmol/l}$, $n = 38$) than in control islets (184




FIG. 5. Concentration-dependency and degree of synchronization of glucose effects on [Ca²⁺]_i in GLP-1R^{+/+} and GLP-1R^{-/-} islets. The glucose concentration of the perfusion medium was decreased stepwise from 15 to 7 mmol/l as indicated. *A*: Mean whole islet [Ca²⁺]_i of 36 GLP-1R^{-/-} and GLP-1R^{+/+} islets. *B* and *C*: Analysis of [Ca²⁺]_i in various sub-regions of islets from GLP-1R^{+/+} (*B*) and GLP-1R^{-/-} mice (*C*). Traces in panels *B* and *C* represent data from 10 of 36 control islets and 36 of 36 knockout islets. As explained in the text, 20 of 36 control islets did not display [Ca²⁺]_i oscillations in the presence of 7 mmol/l glucose. Dotted vertical lines were added to facilitate temporal comparison between traces.

$\pm 4 \text{ nmol/l}$, $n = 38$, $P < 0.01$ for average [Ca²⁺]_i integrated from 1 to 17 min).

We next investigated whether a similar difference in [Ca²⁺]_i is also seen when the islets are stimulated by secretagogues other than glucose. By activating muscarinic receptors, acetylcholine (ACh) exerts complex effects on β -cell [Ca²⁺]_i, which involve both mobilization of cellular Ca²⁺ and stimulation of Ca²⁺ influx (28). Addition of ACh to the medium containing 15 mmol/l glucose rapidly increased [Ca²⁺]_i and transformed the glucose-induced [Ca²⁺]_i oscillations into a sustained elevation that was similar in both types of islets (Fig. 4). Subsequent hyperpolarization of the plasma membrane with diazoxide, a K⁺_{ATP}-channel opener, counteracted the effects of glucose and ACh, and lowered [Ca²⁺]_i to




FIG. 6. Effect of GIP on $[Ca^{2+}]_c$ in GLP-1R $^{+/+}$ and GLP-1R $^{-/-}$ islets. The glucose concentration of the perfusion medium was increased from 3 mmol/l (G3) to 7 mmol/l (G7), and GIP (10 nmol/l) was applied when indicated. The traces are representative (A and B) or mean (C) results from 13 GLP-1R $^{+/+}$ and 12 GLP-1R $^{-/-}$ islets.

near-basal levels. Finally, depolarization with 30 mmol/l K^+ in the presence of diazoxide and absence of ACh rapidly reactivated voltage-dependent Ca^{2+} channels and raised $[Ca^{2+}]_c$ to a sustained plateau that was similar in control and GLP-1R $^{-/-}$ islets (Fig. 4).

The difference in average $[Ca^{2+}]_c$ observed at 15 mmol/l glucose was even more pronounced at 10 and 7 mmol/l (Fig. 5A). When the glucose concentration was raised from 3 to 7 mmol/l, $[Ca^{2+}]_c$ transiently decreased in most control islets whereas it displayed a biphasic rise in GLP-1R $^{-/-}$ islets (Fig. 6). The threshold glucose concentration for induction of a $[Ca^{2+}]_c$ response is thus lower in GLP-1R $^{-/-}$ islets than in control islets.

Digital image analysis was used to test whether the disorganization of glucose-induced $[Ca^{2+}]_c$ oscillations in GLP-1R $^{-/-}$ islets could result from an asynchrony between subregions of the islets. In GLP-1R $^{+/+}$ islets, glucose-induced $[Ca^{2+}]_c$ changes were well synchronized between all subregions even when their amplitude was variable (Fig. 5B). In 10 of 16 GLP-1R $^{+/+}$ islets that displayed $[Ca^{2+}]_c$ oscillations in the presence of 7 mmol/l glucose, some subregions were silent aside from the oscillating ones (Fig. 5B). In contrast to control islets, GLP-1R $^{-/-}$ islets showed poor synchronization of glucose-induced $[Ca^{2+}]_c$ changes between subregions even in 15 mmol/l




FIG. 7. Effect of GLP-1 and GIP on $[Ca^{2+}]_c$ in GLP-1R $^{+/+}$ and GLP-1R $^{-/-}$ islets stimulated with 15 mmol/l glucose. The glucose concentration of the perfusion medium was 15 mmol/l (G15) throughout the experiment. GLP-1 (10 nmol/l) and GIP (10 nmol/l) were applied as indicated. The traces are representative (A and B) or mean (C) results from 48 control and 49 knockout islets.

glucose (Fig. 5C). A sustained $[Ca^{2+}]_c$ response in certain subregions was associated with $[Ca^{2+}]_c$ oscillations in others.

Effect of GLP-1 and GIP on $[Ca^{2+}]_c$. Addition of 10 nmol/l GLP-1 to a medium containing 15 mmol/l glucose induced a small increase in $[Ca^{2+}]_c$ accompanied by a lengthening of the duration of $[Ca^{2+}]_c$ oscillations in 37 of 48 control islets (Fig. 7A), and a sustained plateau in the other islets (not shown). As expected, GLP-1 was ineffective in GLP-1R $^{-/-}$ islets (Fig. 7B). Subsequent addition of GIP raised $[Ca^{2+}]_c$ further and transformed $[Ca^{2+}]_c$ oscillations into a sustained plateau in most control and GLP-1R $^{-/-}$ islets. Average $[Ca^{2+}]_c$ integrated from 24–32 min was not significantly different under these conditions (276 ± 5 nmol/l in 49 GLP-1R $^{-/-}$ islets vs. 259 ± 6 nmol/l in 48 control islets; Fig. 7C). However, when GIP was added to a medium containing 7 mmol/l glucose, it was ineffective in 8 of 13 control islets and induced fast but infrequent $[Ca^{2+}]_c$ oscillations in the other islets (Fig. 6A). By contrast, GIP increased $[Ca^{2+}]_c$ in 12 of 12 GLP-1R $^{-/-}$ islets (Fig. 6B). On average, the GIP-induced $[Ca^{2+}]_c$ rise at 7 mmol/l glucose was much larger in GLP-1R $^{-/-}$ islets than in control islets (Fig. 6C) ($P < 0.01$).

DISCUSSION

It is currently accepted that GLP-1 and GIP are the major incretins in animal models and in man (4,29–31). Furthermore,

it has been proposed that GLP-1 may also be important to keep β -cells—at least part of them—in a glucose-competent state (10,11). We previously suggested that the glucose sensitivity and responsiveness of a pancreatic β -cell population can be explained by a model of heterogeneity among the individual cells (32–34). Since the insulin response of β -cells can be modulated by hormones acting via cAMP (19,20), mouse islets with disrupted GLP-1R signaling (8) are a useful model for studying the role of GLP-1 in the glucose competence of β -cells. The observations presented here demonstrate that, although glucose competence is preserved in perfused GLP-1R^{-/-} islet cells, these islet cells show profound changes in intracellular signal transduction, both in terms of glucose-induced calcium signaling and GIP-induced cAMP production.

We have recently reported (12), and confirm in the present study, that isolated islets from mice homozygous for GLP-1R^{-/-} exhibit a well-preserved insulin-secretory response to glucose. This observation can be interpreted in two ways. First, the presence of GLP-1R is not essential for the secretory response of isolated islets to glucose alone. This possibility is contradicted by our demonstration that the GLP-1R antagonist exendin (9–39)amide inhibited glucose-induced insulin release from control islets during 15-min stimulation either in perfusions or during static incubations. Second, functional GLP-1R are important for maintaining a good secretory response to glucose, but their loss as a result of gene disruption can be compensated for via changes in signaling in β -cells. Our present results showing differences in cAMP accumulation and $[Ca^{2+}]_c$ between GLP-1R^{-/-} and GLP-1R^{+/+} islets support this second hypothesis. cAMP accumulation in the presence of 20 mmol/l glucose was significantly less in GLP-1R^{-/-} islets than in GLP-1R^{+/+} islets, and this difference disappeared completely in the presence of 1 μ mol/l exendin (9–39)amide. This finding suggests that in control islets, even in the absence of ligand, GLP-1R may exhibit a constitutively active role in maintaining the rate of islet cell cAMP production. Consistent with these observations, we previously reported that exendin (9–39)amide lowers basal cAMP accumulation and insulin secretion from rat purified β -cells (14). Others have recently observed reduced basal rate of cAMP formation and decreased insulin secretion from a GLP-1R^{-/-} β -cell line (35). Taken together, these findings suggest that exendin (9–39)amide acts on the “empty” GLP-1R as an inverse agonist. In intact islets of Langerhans and, to a lesser extent, flow-sorted β -cells, paracrine glucagon can accumulate (20). In such conditions, competition between exendin (9–39)amide and glucagon for binding to GLP-1R on β -cells may be an additional mechanism for the effect of exendin (9–39)amide on cAMP accumulation and insulin release (14).

The effect of exendin (9–39)amide on insulin release requires functional GLP-1R and appears cAMP-mediated, because: 1) no effects of exendin (9–39)amide were observed in GLP-1R^{-/-} islets; and 2) in control islets, exendin (9–39)amide did not impair the potentiating effect of 10 nmol/l GIP (Fig. 2B) or Sp-cAMPS (data not shown) on glucose-induced insulin secretion. We do not know why glucose-induced insulin release from control islets incubated for 60 min was not suppressed by exendin (9–39)amide, whereas the effect of a 15-min stimulation was effectively inhibited. One possibility is that GLP-1R might become desensitized

after longer exposure to 1 μ mol/l exendin (9–39)-amide, but the results of the continuous perfusion with exendin (Fig. 1) do not support this idea. Alternatively, 1-h static incubation of control islets might allow sufficient accumulation of paracrine glucagon to compete with exendin (9–39)amide for binding on GLP-1R.

A caveat in the direct comparison of the cAMP data and insulin release data is that the former were collected after incubation with the phosphodiesterase inhibitor IBMX, thus allowing to detect differences in cAMP production, while insulin release was measured in the absence of IBMX. In control experiments, in which we added 250 μ mol/l IBMX to the 15-min statically incubated islets, the same trends on insulin release after adding GLP-1 and GIP were observed as in the absence of IBMX, but the responses were much smaller, largely because the effect of glucose alone was very high (data not shown). It should not be overlooked, therefore, that the difference in “basal” cAMP concentration of control and knockout β -cells may be smaller in vivo than what was measured in the present conditions, first because the in vitro measurements required addition of a phosphodiesterase inhibitor and, second, because the contribution of non- β -cells to islet cAMP in this model is not known.

Assuming that the acute suppression of glucose-induced insulin release proceeds via lowering of intracellular cAMP accumulation, our current and previous (14) data suggest a role for a sufficiently high level of basal cAMP production in the insulin-secretory response of islets to high glucose. Such a conclusion is not supported, however, by a recent study (36) showing that disruption of protein kinase A anchoring in rat islets and RINm5F cells by introduction of anchoring peptides or expression of soluble A-kinase-anchoring proteins blocked GLP-1-stimulated but not glucose-induced insulin release. The experimental designs (the species used, acute versus chronic disruption of islet cAMP signaling) in the study by Lester et al. (36) and the present study are quite different, so that it seems difficult to directly compare results.

Because glucose-induced insulin secretion from GLP-1R^{-/-} islets is well preserved (12 and this study) despite lower cAMP levels, it can be postulated that one or several mechanisms compensate for the disruption of the GLP-1R signaling. As a first possibility, we considered compensation via an increase in the sensitivity to cAMP, e.g., by upregulating catalytic subunits, downregulating regulatory subunits of cAMP-dependent protein kinase or changes in the expression of targets of cAMP-dependent phosphorylation. This possibility was assessed by measuring insulin secretion at 20 mmol/l glucose in the presence of different concentrations of Sp-cAMPS, a potent cAMP analog that is taken up by intact cells (37). The absence of enhanced sensitivity to Sp-cAMPS suggests that increased cAMP sensitivity, particularly at higher levels of intracellular cAMP, is not a generalizable feature of GLP-1R^{-/-} islets. Second, because GLP-1 increases $[Ca^{2+}]_c$ in beta-cells (38) and mobilizes intracellular Ca^{2+} in β TC3-cells (39), we evaluated whether compensating mechanisms in islets of GLP-1R^{-/-} mice could involve changes in $[Ca^{2+}]_c$. A consistent feature of GLP-1R^{-/-} islets was a higher average $[Ca^{2+}]_c$ in the presence of a stimulating glucose concentration. An increased glucose-induced $[Ca^{2+}]_c$ response is unexpected in the context of a decrease in islet cAMP and the knowledge that

cAMP has a slight positive effect on Ca^{2+} -channel activity in β -cells (21,22). Because the islets isolated from both mouse strains responded identically to high K^+ in the presence of diazoxide, it seems unlikely that the higher $[\text{Ca}^{2+}]_c$ response to glucose in $\text{GLP-1R}^{-/-}$ islets is the result of an increased number of voltage-dependent Ca^{2+} channels. Furthermore, ACh had similar effects on $[\text{Ca}^{2+}]_c$ in both types of islets, indicating that inositol-1,4,5-triphosphate (IP_3)-mediated calcium mobilization from intracellular stores has not been changed substantially in $\text{GLP-1R}^{-/-}$ β -cells. It is conceivable that a chronic decrease in cAMP production has effects on Ca^{2+} channel function that are different from those measured in acute in vitro experiments.

So, how might this change in glucose-signaling be achieved? One possibility, to be examined by future experiments, is that higher $[\text{Ca}^{2+}]_c$ in $\text{GLP-1R}^{-/-}$ β -cells at stimulatory glucose concentrations results from an increase in glucose metabolism. This idea is substantiated by the observation that the threshold glucose concentration for induction of a $[\text{Ca}^{2+}]_c$ response was lower in $\text{GLP-1R}^{-/-}$ than in $\text{GLP-1R}^{+/+}$ islets. Increased metabolism would not only affect $[\text{Ca}^{2+}]_c$ through the K^+ -ATP dependent pathway, but also potentiate insulin secretion through the K^+ -ATP channel-independent pathway (40). Interestingly, we observed signs of asynchronism of $[\text{Ca}^{2+}]_c$ oscillations in response to glucose between subregions in islets of $\text{GLP-1R}^{-/-}$ mice. Among several mechanisms that could explain this phenomenon, a difference in metabolic and electric coupling between islet β -cells should be considered. Such coupling has been shown previously to depend on gap junctions between islet cells (41,42). In other tissues it has been shown that gap junctional communication is increased by raising the intracellular cAMP concentration (43,44). Therefore, the low cAMP content in $\text{GLP-1R}^{-/-}$ islets may be insufficient to maintain adequate coupling between large numbers of islet cells. Taken together, our data seem to indicate that multiple abnormalities in terms of $[\text{Ca}^{2+}]_c$ and cAMP exist in $\text{GLP-1R}^{-/-}$ islets and that the larger $[\text{Ca}^{2+}]_c$ response to glucose of $\text{GLP-1R}^{-/-}$ β -cells compensates, at least in terms of glucose-induced insulin release, for lower cAMP levels.

We have previously reported that $\text{GLP-1R}^{-/-}$ mice exhibit upregulation of the GIP system, both in terms of GIP plasma levels and in terms of GIP potentiation of glucose-induced insulin release (13). The present study extends these findings by demonstrating a higher responsiveness of $\text{GLP-1R}^{-/-}$ islets to GIP at the level of islet cAMP accumulation (Fig. 2A). This hyperresponsiveness could result from upregulation of the number of GIP receptors or increased signaling following activation of the GIP receptor. Consistent with the cAMP data are the significantly larger effects of 10 nmol/l GIP on 20 mmol/l glucose-induced insulin release (Fig. 2B) and on 7 mmol/l glucose-induced $[\text{Ca}^{2+}]_c$ (Fig. 7) in $\text{GLP-1R}^{-/-}$ islets.

In summary, the data presented in this article suggest that, despite the well-preserved insulin secretory response to glucose of $\text{GLP-1R}^{-/-}$ mice in vitro, $\text{GLP-1R}^{-/-}$ islets exhibit profound changes in basal and stimulated intracellular signaling at the level of both cAMP and $[\text{Ca}^{2+}]_c$. It remains to be shown to what extent upregulation of GIP-R-induced cAMP accumulation and the larger $[\text{Ca}^{2+}]_c$ response to glucose in $\text{GLP-1R}^{-/-}$ β -cells are the pivotal mediators of in vivo compensatory adaptation to the genetic loss of β -cell GLP-1R signaling.

ACKNOWLEDGMENTS

This study was supported by the Belgian Funds for Scientific Research (Fonds voor Wetenschappelijk Onderzoek [FWO] Vlaanderen Grants 7.0004.96 and G.0130.99 to F.S. and Fonds National de la Recherche Scientifique [FNRS] Grants 3.4552.98 to J.C.H. and 2.4614.99 to P.G.), the Belgian Program on Interuniversity Poles of Attraction (IUAP IV) initiated by the Belgian State, and the Juvenile Diabetes Foundation (D.J.D.).

We are grateful to Prof. J. Vandekerckhove (Department of Physiological Chemistry, University of Ghent) for providing synthetic exendin (9–39)amide. We thank V. Berger for technical assistance.

REFERENCES

1. Elrick H, Stummel L, Illad CJ Jr, Arai Y: Plasma insulin response to oral and intravenous glucose administration. *J Clin Endocrinol Metab* 24:1076–1082, 1964
2. Creutzfeldt W: The incretin concept today. *Diabetologia* 16:75–85, 1979
3. Nauck M, Stockmann F, Ebert R, Creutzfeldt W: Reduced incretin effect in type 2 (non-insulin-dependent) diabetes. *Diabetologia* 29:46–52, 1986
4. Drucker DJ: Glucagon-like peptides. *Diabetes* 47:159–169, 1998
5. Nauck MA, Heimesaat MM, Orskov C, Holst JJ, Ebert R, Creutzfeldt W: Preserved incretin activity of glucagon-like peptide 1 (7–36 amide) but not of synthetic human gastric inhibitory polypeptide in patients with type-2 diabetes mellitus. *J Clin Invest* 91:301–307, 1993
6. Ebert R, Illmer K, Creutzfeldt W: Release of gastric inhibitory polypeptide (GIP) by intraduodenal acidification in rats and humans and abolishment of the incretin effect of acid by GIP-antiserum in rats. *Gastroenterology* 76:515–523, 1979
7. Göke R, Fehmman HC, Linn T, Schmidt H, Krause M, Eng J, Göke B: Exendin-4 is a high potency agonist and truncated exendin (9–39)amide an antagonist at the glucagon-like peptide 1 (7–36)amide receptor of insulin-secreting β -cells. *J Biol Chem* 268:19650–19655, 1993
8. Scrocchi LA, Brown TJ, Maclusky N, Brubaker PL, Auerbach AB, Joyner AL, Drucker DJ: Glucose intolerance but normal satiety in mice with a null mutation in the glucagon-like peptide 1 receptor gene. *Nat Med* 2:1254–1258, 1996
9. Gutniak M, Orskov C, Holst JJ, Ahrén B, Efendic S: Antidiabetogenic effects of glucagon-like peptide-1 (7–36)amide in normal subjects and patients with diabetes mellitus. *New Engl J Med* 326:1316–1322, 1992
10. Holz GG IV, Kühtreiber WM, Habener JF: Pancreatic beta-cells are rendered glucose-competent by the insulinotropic hormone glucagon-like peptide-1 (7–37). *Nature* 361:362–365, 1993
11. Montrose-Rafizadeh C, Wang Y, Janczewski AM, Henderson TE, Egan JM: Overexpression of glucagon-like peptide-1 receptor in an insulin-secreting cell line enhances glucose responsiveness. *Mol Cell Endocrinol* 130:109–117, 1997
12. Flamez D, Van Breusegem A, Scrocchi LA, Quartier E, Pipeleers D, Drucker DJ, Schuit F: Mouse pancreatic β -cells exhibit preserved glucose competence after disruption of the glucagon-like peptide-1 receptor gene. *Diabetes* 47:646–652, 1998
13. Pederson RA, Sathkumarajah M, McIntosh CHS, Scrocchi LA, Flamez D, Schuit F, Drucker DJ, Wheeler MB: Enhanced GIP secretion and insulinotropic action in glucagon-like peptide receptor $-/-$ mice. *Diabetes* 47:1046–1052, 1998
14. Moens K, Flamez D, Van Schravendijck C, Ling Z, Pipeleers D, Schuit F: Dual glucagon recognition by pancreatic β -cells via glucagon and glucagon-like peptide 1 receptors. *Diabetes* 47:66–72, 1998
15. Moens K, Heimberg H, Flamez D, Huypens P, Quartier E, Ling Z, Pipeleers D, Gremlich S, Thorens B, Schuit F: Expression and functional activity of glucagon, glucagon-like peptide 1, and glucose-dependent insulinotropic peptide receptors in rat pancreatic islet cells. *Diabetes* 45:257–261, 1996
16. Drucker DJ, Philippe J, Mojsos S, Chick WL, Habener JF: Glucagon-like peptide I stimulates insulin gene expression and increases cyclic AMP levels in a rat islet cell line. *Proc Natl Acad Sci U S A* 84:3434–3438, 1987
17. Fehmman HC, Janssen M, Göke B: Interaction of glucagon-like peptide-I (GLP-1) and galanin in insulin (beta TC-1)- and somatostatin (RIN T3)-secreting cells and evidence that both peptides have no receptors on glucagon (INRIG9)-secreting cells. *Acta Diabetologica* 32:176–181, 1995
18. Gromada J, Holst JJ, Rorsman P: Cellular regulation of islet hormone secretion by the incretin hormone glucagon-like peptide-1. *Pflügers Arch Eur J Physiol* 435:583–594, 1998
19. Pipeleers DG, Schuit FC, In't Veld P, Maes E, Hooghe-Peters EL, Van De Winkel M, Gepts W: Interplay of nutrients and hormones in the regulation of

- insulin release. *Endocrinology* 117:824-833, 1985
20. Schuit FC, Pipeleers DG: Regulation of adenosine 3',5'-monophosphate levels in the pancreatic B cell. *Endocrinology* 117:834-840, 1985
 21. Ämmälä C, Ashcroft FM, Rorsman P: Calcium-independent potentiation of insulin release by cyclic AMP in single β -cells. *Nature* 363:356-358, 1993
 22. Henquin JC, Meissner HP: The ionic, electrical, and secretory effects of endogenous cyclic adenosine monophosphate in mouse pancreatic B cells: studies with forskolin. *Endocrinology* 115:1125-1134, 1984
 23. Gilon P, Shepherd RM, Henquin JC: Oscillations of secretion driven by oscillations of cytoplasmic Ca²⁺ as evidenced in single pancreatic islets. *J Biol Chem* 268:22265-22268, 1993
 24. Pralong WF, Bartley C, Wollheim CB: Single islet β -cell stimulation by nutrients: relationship between pyridine nucleotides, cytosolic Ca²⁺ and secretion. *EMBO J* 9:53-60, 1990
 25. Ling Z, Kiekens R, Mahler T, Schuit FC, Pipeleers-Marichal M, Sener A, Kloppel G, Malaisse WJ, Pipeleers DG: Effects of chronically elevated glucose levels on the functional properties of rat pancreatic beta-cells. *Diabetes* 45:1774-82, 1996
 26. Gilon P, Henquin JC: Influence of membrane potential changes on cytoplasmic Ca²⁺ concentration in an electrically excitable cell, the insulin-secreting pancreatic B-cell. *J Biol Chem* 267:20713-20720, 1992
 27. Renström E, Eliasson L, Rorsman P: Protein kinase-dependent and -independent stimulation of exocytosis by cAMP in mouse pancreatic B-cells. *J Physiol* 502:105-118, 1997
 28. Gilon P, Nenquin M, Henquin JC: Muscarinic stimulation exerts both stimulatory and inhibitory effects on the concentration of cytoplasmic Ca²⁺ in the electrically excitable pancreatic β -cell. *Biochem J* 311:259-267, 1995
 29. Thorens B: Expression cloning of the pancreatic β -cell receptor for the glucagon-like peptide 1. *Proc Natl Acad Sci U S A* 89:8641-8645, 1992
 30. Tseng CC, Kieffer TJ, Jarboe LA, Usdin TB, Wolfe MM: Postprandial stimulation of insulin release by glucose-dependent insulinotropic polypeptide (GIP). *J Clin Invest* 98:2440-2445, 1996
 31. Wang Z, Wang RM, Owji AA, Smith DM, Ghatei MA, Bloom SR: Glucagon-like peptide-1 is a physiological incretin in rat. *J Clin Invest* 95:417-421, 1995
 32. Schuit FC, In't Veld PA, Pipeleers DG: Glucose stimulates proinsulin biosynthesis by a dose-dependent recruitment of pancreatic beta cells. *Proc Natl Acad Sci U S A* 85:3865-3869, 1988
 33. Kiekens R, In't Veld P, Mahler T, Schuit F, Van De Winkel M, Pipeleers D: Differences in glucose recognition by individual rat pancreatic B cells are associated with intercellular differences in glucose-induced biosynthetic activity. *J Clin Invest* 89:117-125, 1992
 34. Heimberg H, De Vos A, Vandercammen A, Van Schaftingen E, Pipeleers D, Schuit F: Heterogeneity in glucose sensitivity among pancreatic β -cells is correlated to differences in glucose phosphorylation rather than in glucose transport. *EMBO J* 12:2873-2879, 1993
 35. Serre V, Dolci W, Schaerer E, Scrocchi L, Drucker D, Efrat S, Thorens B: Exendin (9-39)amide is an inverse agonist of the murine GLP-1 receptor: implications for basal intracellular cyclic adenosine 3', 5'-monophosphate levels and β -cell glucose competence. *Endocrinology* 139:4448-4454, 1998
 36. Lester LB, Langeberg LK, Scott JD: Anchoring of protein kinase A facilitates hormone-mediated insulin secretion. *Proc Natl Acad Sci U S A* 94:14942-14947, 1997
 37. Sammak PJ, Adams SR, Harootyan AR, Schliwa M, Tsien RY: Intracellular cAMP not calcium, determines the direction of vesicle movement in melanophores: direct measurement by fluorescence ratio imaging. *J Cell Biol* 117:57-72, 1992
 38. Holz GG IV, Leech CA, Habener JF: Activation of a cAMP-regulated Ca²⁺-signaling pathway in pancreatic β -cells by the insulinotropic hormone GLP-1. *J Biol Chem* 270:17749-17757, 1995
 39. Gromada J, Rorsman P: Molecular mechanism underlying GLP-1 induced calcium mobilization from internal stores in insulin-secreting β -TC3 cells. *Acta Physiol Scand* 157:349-351, 1996
 40. Gembal M, Gilon P, Henquin JC: Evidence that glucose can control insulin release independently from its action on ATP-sensitive K⁺ channels in mouse β -cells. *J Clin Invest* 89:1288-1295, 1992
 41. Halban PA, Wollheim CB, Blondel B, Meda P, Niesor EN, Mintz DH: The possible importance of contact between pancreatic islet cells for the control of insulin release. *Endocrinology* 111:86-94, 1982
 42. Meda P, Michaels RL, Halban PA, Orci L, Sheridan JD: In vivo modulation of gap junctions and dye coupling between B-cells of the intact pancreatic islet. *Diabetes* 32:858-868, 1983
 43. Saez JC, Spray DC, Nairn AC, Hertzberg EL, Greengard P, Bennett MVL: cAMP increases junctional conductance and stimulates phosphorylation of the 27-kDa principal gap junction polypeptide. *Proc Natl Acad Sci U S A* 83:2473-2477, 1986
 44. Murray SA, Fletcher WH: Hormone-induced intercellular signal transfer dissociates cyclic AMP-dependent protein kinase. *J Cell Biol* 98:1710-1719, 1984